




Alexandre Baffet  
Team Leader  
alexandre.baffet@curie.fr  
Tel: +33 1 56 24 63 60

**The neocortex is the center for higher brain functions, such as perception, decision-making and language. Our group focuses on the mechanisms governing neocortex development, with a strong interest on the role and regulation of the neural stem cells.**

Cortical development occurs by proliferation of neural stem cells and migration of newborn neurons over substantial distances. In mice, neuroepithelial cells differentiate into apical radial glial cells (aRG), which act as the stem cells of the developing neocortex and give rise to all neocortical neurons, most glial cells and to the adult stem cells. aRG cells are highly elongated, extending a basal process all the way to the pial surface of the developing brain and an apical process that remains in contact with the ventricular surface (**Figure. 1**). They undergo fascinating cell cycle-dependent nuclear oscillations, a process known as Interkinetic Nuclear Migration (INM). We previously described the mechanism for G2 apical nuclear migration, which involves a Cdk1-triggered recruitment of the dynein motor to the nuclear pore complex (Hu et al., 2013; Baffet et al., 2015). aRG cells can divide asymmetrically to produce one aRG cell and one intermediate progenitor that will divide to produce two neurons.



**Figure 1. Mouse and human neocortex development.**

*In mouse, aRG cells are the major neural stem cells. They are highly elongated, with an apical process in contact with the ventricular surface and a basal process connected to the pial surface of the brain. In human, the bRG cells (orange) have lost connection to the apical surface and are localized basally, in the highly enlarged OSVZ. I.P.: Intermediate Progenitor.*


Although the ventricular zone (VZ), where aRG

cells are located, is the major neurogenic region in mice, the situation is very different in humans where 85% of mitoses occur in the sub-ventricular zone (SVZ). As a consequence, this region is massively thicker and becomes subdivided into two morphologically different regions, the inner SVZ (ISVZ) and the outer SVZ (OSVZ). This is largely due to a second type of neural stem cells, called the basal Radial Glial (bRG) cells (also known as oRG cells) (**Figure 1**). bRG cells derive from aRG cells and share several characteristics with these cells, such as the extension of a basal process, the expression of a similar set of transcription factors and, most importantly, the capacity to self-renew. bRG cells are very rare in the smooth mouse brain (lissencephalic) and very abundant in the folded human brain (gyrencephalic) and are thought to be fundamental players of the massive size increase of the human brain.

## Our group currently focuses on three main topics:

### 1/ Mechanisms of polarized trafficking in RG cells


In the highly elongated aRG and bRG cells, long-range polarized trafficking occurs to deliver critical molecules, including surface receptors or secreted factors. To investigate these processes, we have recently developed a method for high resolution subcellular live imaging within thick embryonic brain slices. This approach allows to resolve individual fast-moving structures, such as RAB6+ positive vesicles while they are transported in the apical or basal process (**Figure 2**).



**Figure 2. Subcellular live imaging in RG cells in situ.**

**A.** Experimental set-up for mouse embryonic brain in utero electroporation and brain slice preparation. **B.** Live imaging of GFP-RAB6 in mouse radial glial cell in situ.

Using this method, we have identified the organization of the microtubule cytoskeleton in RG cells (Coquand et al., 2020). We showed that, while microtubules in the apical process of aRG cells in uniformly oriented basally, microtubules in the basal process display a mixed polarity, reminiscent of the mammalian dendrite (**Figure 3**). We furthermore showed that this acentrosomal microtubule network is organized from CAMSAP+ varicosities of the basal process, an organization conserved in human bRG cells. We are currently initiating a project aimed at identifying the factors that control this acentrosomal bipolar microtubule network organization.



**Figure 3. Microtubule organization in RG cells in situ.**


Microtubules in the apical process have a unipolar basal orientation, while microtubules in the basal process are bipolar and are organized from varicosities.

Our second project aims at identifying the mechanism for post-Golgi apical trafficking in epithelial cells, using aRG cells as a model system (Brault et al, *in preparation*). Using our live imaging method, as well as a newly generated RAB6A/B double knock-out mouse, we have identified the core machinery that drives post-Golgi transport of the Crumbs apical determinant to the apical surface of mouse aRG cells. We show that impairment of this pathways leads to a delamination of aRG cells that now divide basally, adopting a bRG-like identity. We have now begun a novel project aimed at identifying the mechanisms for basal secretion in RG cells.

## 2/ Human neurogenesis

To image human bRG cells, we have established the generation of iPS cell-derived human cerebral organoids. We observe the appearance of bRG cells after 5 weeks of culture, which

gradually increases in the following weeks. While organoids usually display mild amounts of bRG cells, our adapted protocol leads to the formation of a large OSVZ containing a very abundant bRG cell population (**Figure 4A**). In parallel, we have developed the culture and imaging of human fetal brain tissue, obtained through collaborations with two Parisian hospitals (**Figure 4B**). Because of the dynamic nature of the processes we investigate, we have developed methods for quantitative live imaging of human fetal tissue and cerebral organoids. Tissues are sliced and put into culture for 7 days. Plasmids encoding fluorescent reporters or knockdown constructs are delivered using retroviral infection and slices are live imaged using a CSU-W1 Spinning wide microscope for 24 to 48 hours (**Figure 4C, D**). Finally, we have developed a protocol to generate *in vitro* cultures of human RG cells, obtained from human fetal tissue or cerebral organoids. We have developed assays based on microfabrication techniques allowing to align them on micropatterns of adhesion, in order to manipulate their shape (**Figure 4E, F**).



**Figure 4. Methods for the investigation of human bRG cells.**

**A.** Human cerebral organoid stained for the RG cell marker Sox2, the intermediate progenitor marker Tbr2 and the neuronal marker NeuN. **B.** Human fetal brain slice stained for the RG cell marker Sox2

and the mitotic RG cell marker Phospho-Vimentin (P-Vim). **C.** Schematic representation of brain slice infection and *in vitro* RG cell culture from fetal brain tissue biopsy. **D.** Live imaging of migrating and dividing bRG cell in human fetal cortex. **E.** *In vitro* cultivated human RG cells. **F.** A RG cell aligned on an elongated fibronectin pattern.

We currently pursue two main projects that focus on human bRG cells. First, we investigate how fate decisions are made in these cells. We have developed a live-fix correlative microscopy method, that allows us to live image dividing bRG cells *in situ*, and to identify the fate of the daughter cells. This approach has allowed us to generate a map of bRG cell division modes. Remarkably, this map is highly conserved between human fetal tissue and cerebral organoids (Coquand et al, *in preparation*). We are now using this knowledge to identify the mechanisms controlling asymmetric cell division, whereby one bRG cells divides into one self-renewing bRG cell and one differentiating cell. In a second project, we are investigating the cytoskeletal-based mechanisms that drive the migration of bRG cells into the developing human neocortex.

### 3/ Brain Malformations

Alterations in RG cells can lead to a variety of cerebral malformations, which can have genetic or environmental causes (including viral infection). We developed an assay to infect brain slices with flaviviruses and demonstrated that the microcephaly-causing Zika virus (but not other closely-related flaviviruses) specifically infect RG cells and lead to apoptotic cell death (Brault et al., 2016).

In another project, our collaborators have identified mutations in the WDR81 gene that causes extreme microcephaly in newborn children. Using a newly generated WDR81 KO mouse model as well as patient-derived cells, we demonstrated that reduced brain size is specifically due to reduced proliferation rates of RG cells. We showed that WDR81 regulates endosomal trafficking of EGFR, and that loss of function leads to reduced MAP kinase pathway activation (Carpentieri et al., 2020).

Finally, we use patient-derived iPS cells and cerebral organoids to model how mutations can affect human brain development. In particular, we focus on mutations in the DHC1H1 gene, which codes for the dynein heavy chain. Strikingly, our collaborators have identified clusters of mutations that can lead either to cortical malformations, or to motor neuron degeneration (Spinal Muscular Atrophy). We investigate how these different types of mutations differentially affect dynein-based process in the developing cerebral tissue (Farcy et al, *in preparation*).

## Key publications

### Year of publication 2020

---

Jacopo A. Carpentieri, Amandine Di Cicco, David Andreau, Laurence Del Maestro, Fatima El

Marjou, Laure Coquand, Jean-Baptiste Brault, Nadia Bahi-Buisson, Alexandre D. Baffet (2020 Aug 17)

**Endosomal trafficking defects alter neural progenitor proliferation and cause microcephaly**

*BioRxiv* : [DOI : doi.org/10.1101/2020.08.17.254037](https://doi.org/10.1101/2020.08.17.254037)

L. Coquand, G.S. Victoria, A. Tata, J.B. Brault, F. Guimiot, V. Fraissier, A. D. Baffet (2020 Mar 16)

**A dendritic-like microtubule network is organized from swellings of the basal fiber in neural progenitors**

*BioRxiv* : [DOI : 10.1101/2020.03.16.993295](https://doi.org/10.1101/2020.03.16.993295)

**Year of publication 2016**

---

Jean-Baptiste Brault, Cécile Khou, Justine Basset, Laure Coquand, Vincent Fraissier, Marie-Pascale Frenkiel, Bruno Goud, Jean-Claude Manuguerra, Nathalie Pardigon, Alexandre D Baffet (2016 Jul 26)

**Comparative Analysis Between Flaviviruses Reveals Specific Neural Stem Cell Tropism for Zika Virus in the Mouse Developing Neocortex.**

*EBioMedicine* : [DOI : S2352-3964\(16\)30323-1](https://doi.org/10.1016/S2352-3964(16)30323-1)

Alexandre D Baffet, Aurélie Carabalona, Tiago J Dantas, David D Doobin, Daniel J Hu, Richard B Vallee (2016 Jan 23)

**Cellular and subcellular imaging of motor protein-based behavior in embryonic rat brain.**

*Methods in cell biology* : 349-63 : [DOI : 10.1016/bs.mcb.2015.06.013](https://doi.org/10.1016/bs.mcb.2015.06.013)

**Year of publication 2015**

---

Alexandre D Baffet, Daniel J Hu, Richard B Vallee (2015 Jun 9)

**Cdk1 Activates Pre-mitotic Nuclear Envelope Dynein Recruitment and Apical Nuclear Migration in Neural Stem Cells.**

*Developmental cell* : 703-16 : [DOI : 10.1016/j.devcel.2015.04.022](https://doi.org/10.1016/j.devcel.2015.04.022)

**Year of publication 2014**

---

Alexandre D Baffet (2014 Jan 30)

**[Nuclear migration in neuronal progenitors: when the brain plays yo-yo].**

*Médecine sciences : M/S* : 30-2 : [DOI : 10.1051/medsci/20143001009](https://doi.org/10.1051/medsci/20143001009)